

South Dakota State University

Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange

Theses and Dissertations

1937

The Effect of Cultivation on Certain Chemical and Physical Properties of Some South Dakota Soils

Oscar E. Olson

Follow this and additional works at: <http://openprairie.sdstate.edu/etd>



Part of the [Agronomy and Crop Sciences Commons](#), and the [Soil Science Commons](#)

Recommended Citation

Olson, Oscar E., "The Effect of Cultivation on Certain Chemical and Physical Properties of Some South Dakota Soils" (1937). *Theses and Dissertations*. 1967.

<http://openprairie.sdstate.edu/etd/1967>

This Thesis - Open Access is brought to you for free and open access by Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. It has been accepted for inclusion in Theses and Dissertations by an authorized administrator of Open PRAIRIE: Open Public Research Access Institutional Repository and Information Exchange. For more information, please contact michael.biondo@sdstate.edu.

TABLE OF CONTENTS

**THE EFFECT OF CULTIVATION
ON CERTAIN CHEMICAL AND PHYSICAL PROPERTIES
OF SOME SOUTH DAKOTA SOILS**

I. Acknowledgments	1
II. Introduction	2
III. Review of Literature	3
IV. Purpose of Investigation	6
V. Plan of Investigation	7
A. Thesis Submitted to the Faculty of South Dakota State College by Oscar E. Olson	7
B. Description of Soil Series	10
VI. Discussion of Results	13
A. Physical Analysis	13
1. Methods of Analysis	13
2. Aggregate analysis	13
3. Structural analysis	13
4. Mechanical analysis	15
B. Chemical Analysis	20
1. Methods of analysis	20
2. Loss on ignition	20
3. Organic matter	21
4. Humus	21
5. Nitrogen	21
6. Phosphorus	22
7. Soil reaction	24
8. Base exchange capacity	25
VII. Summary	26
VIII. Literature cited	27

THIS BOOK DOES
NOT CIRCULATE

SOUTH DAKOTA
STATE COLLEGE LIBRARY

In Partial Fulfillment of the Requirements for the Degree
of Master of Science

June, 1937

TABLE OF CONTENTS

	Page
I. Acknowledgements	
II. Introduction	1
III. Review of Literature	3
IV. Purpose of Investigation	6
V. Plan of Investigation	7
A. Collection of Materials	7
1. Description and location of samples	9
B. Description of Soil Series	10
VI. Discussion of Results	13
A. Physical Analysis	13
1. Methods of Analysis	13
2. Aggregate analysis	13
3. Structural stability	15
4. Mechanical analysis	25
B. Chemical Analysis	28
1. Methods of analysis	28
2. Loss on ignition	28
3. Organic matter	30
4. Humus	32
5. Nitrogen	33
6. Phosphorus	36
7. Soil reaction	38
8. Base exchange capacity	40
VII. Summary	
VIII. Literature cited	


6 June 37 - Auction - 9.

ACKNOWLEDGEMENT

The writer wishes to express his sincere gratitude to Professor Leo F. Puhr for his continued guidance and assistance in conducting this study. Appreciation is also expressed for the cooperation of Dr. A. N. Hume and Professor J. G. Hutton of the Agronomy Department.

Figure 1

PRINCIPAL SOIL AREAS IN SOUTH DAKOTA



LEGEND

- | | |
|--------------------------|--------------------|
| A--Missouri Bottom Soils | M--Morton Soils |
| B--Bad Land Soils | P--Pierre Soils |
| C--Cheyenne Loam Soils | R--Rosebud Soils |
| D--Dune Sand | S--Smithwick Soils |
| G--Glacial Soils | W--Loessial Soils |
| L--Lacustrine Soils | BH--Black Hills |

(From map by J. G. Hutton,
in charge of South Dakota
Soil Survey.)

INTRODUCTION

Soil nutrient losses have followed closely in the paths of advancing frontiers in agriculture. Remedies for depletion as the result of overcropping were sought by George Washington, whose interest and investigation of soil nutrient losses probably represent the earliest attention given to land exploitation in the United States, which at the present time has become a problem of national concern.

Soil losses have been attributed to a number of causes, among which are cropping, erosion, leaching, burning, and rapid oxidation of organic matter in the soil. Associated with soil nutrient losses, certain significant physical and chemical changes may occur in the soil.

The soils of South Dakota lie entirely in the area commonly designated as the Pedocal group. The characteristic feature of the Pedocal group is a horizon of calcium carbonate accumulation in the soil profile which indicates that these soils have developed in an environment of limited rainfall.

It has been generally observed that in regions of limited rainfall the agricultural practices which have prevailed since virgin soils were cultivated, have resulted in the loss of considerable of the original organic matter and nitrogen. The cropping systems commonly practiced have not included the planting of a legume to restore the nitrogen and organic

matter losses due to cultivation and much of the organic matter of the non-leguminous crop residues are partially lost by feeding to livestock, while some are totally lost by burning.

Associated with the loss of organic matter certain physical changes may occur in the soil which have a bearing upon soil structure. Diminution of organic matter of soils, and the probable subsequent change in soil structure, have been regarded as contributory factors in the severity of wind erosion, and to some extent water erosion in the Great Plains area.

The present study aims to make some contribution towards an understanding of the effect of one factor, namely, the cultivation of South Dakota soils, upon (1) the nutrient losses as determined by changes in chemical composition of the soil, and upon (2) the physical properties of the soil.

Result of cultivation. E. E. Viets, of Oklahoma, observed that in native soils cultivation has resulted in loss in nitrogen, no decrease in fertility, and the removal of the upper one-fourth to one-half inch of soil erosion. In native soils and in soils which have been removed, and the fertility has been reduced five to ten per cent after ten years of cultivation.

...the native soils that have been in use for several years."

REVIEW OF LITERATURE

In the report of the eleventh annual meeting of the American Soil Survey Association (2) the observations of several field men on the cultural changes in soils are summarized. G. W. Conrey states that in certain soils (Superior clay in Wisconsin and Crosby silty clay in Ohio) cultivation has changed the texture of the soil to such an extent that a fence line may separate two soils so radically different as to be very confusing at times. W. O. Veatch of Michigan found mixing of the horizons and either gains or losses, depending upon factors such as the type of soil, methods of farming, etc., in the nitrogen, phosphorus, and lime content. W. J. Geib states that in Wisconsin soils the most marked change due to cultivation is the loss in organic matter. A. H. Joel of Saskatchewan found a decrease in crude organic matter and a breakdown of structure as the result of cultivation. F. G. Fitzpatrick of Oklahoma observed that in mature soils cultivation has resulted in loss in nitrogen, no decrease in fertility, and the removal of the upper one-fourth to one-half inch of soil erosion. In immature soils one to four inches of soil has been removed, and the fertility has been reduced five to ten per cent after ten years of cultivation.

Bradfield (8) states, "The best structure is usually found in virgin soils that have been in sod for several years."

A slight increase in the sand content of cultivated Southern High Plains soils over virgin soils of the same area, due to cropping and wind erosion, was reported by Daniel (10).

Daniel and Langham (11) found a decrease in organic matter and total nitrogen in Oklahoma Panhandle soils, due to cultivation.

DeTurk (13) found a migration of colloids from the surface soils downward as the result of cultivation of Illinois soils. He found also that crop rotations commonly practiced induce a shift in organic matter to lower levels than those of the virgin state. He states that, "Plant residues return phosphorus mainly to the surface soil, while root systems remove it from the surface and also from greater depths."

DeTurk, Bauer, and Smith (14) estimated that, based on crop yield, a loss of six pounds of phosphorus per year from Illinois soils, especially the top 6-2/3 inches, occurred. They found no pronounced differences in the total amounts of calcium, magnesium, or sulfur due to cropping or cultivation, except for a gain in calcium in the surface soil of treated plots where calcium was added both as lime and as the phosphate.

With the exception of nitrogen, Dorman (15) reports that in Michigan soils a comparison of analyses of cropped and virgin soils showed no consistent differences. He found that in lighter textured soils phosphorus was decreased by cropping and in heavier soils increased. With reference to the effect of cultivation and cropping, no change in pH was found, and

only small differences in exchangeable hydrogen were noted; both increases and decreases in base exchange capacity were found; with one exception, readily available phosphorus was greater in the virgin soil; and in the cropped soils the greatest response to nitrate fertilizers resulted.

Miller (17) studied the rate of solution of virgin and cultivated soils, using the freezing point method, and found that virgin soils possess the power of giving up soluble material at a greater rate than the corresponding cropped soils. He also found that subsoils from depleted areas had as great a solubility as subsoils from the corresponding virgin soils, indicating that most crop plants feed primarily in the surface or plowed stratum of the soil.

Shedd (19), in Kentucky soils, found that in most cases a loss in calcium was found as the result of cultivation. Many soils were found to be so low in calcium that their deficiency in this constituent becomes as important as deficiencies in phosphorus and nitrogen.

Swanson (2) found in Kansas soils that from 22.6% to 43.5% of the nitrogen and from 23.3% to 51.3% of the organic matter, based on the virgin soil content, was lost due to cultivation.

Swanson and Miller (21) in a study of Kansas soils, report losses in sulfur from the surface soils greater than the amount removed in crops, and concluded that sulfonation and leaching were the cause.

The findings of the majority of the soil investigators which are pertinent to the interests of the present study,

seem to indicate a general loss of nitrogen, calcium, and phosphorus in cultivated soils when compared with virgin soils. These investigators point to some changes in soil texture and structure caused by cultivation. Base exchange capacity and rate of solution of soil nutrients was found to be lower in cultivated soils. Reduced fertility and erosion were generally observed. Probably the most pronounced loss due to cultivation was the loss of organic matter.

PURPOSE OF INVESTIGATION

From the agricultural viewpoint any chemical and physical changes in the soil brought about by cultivation which have a bearing on soil erosion and soil fertility are highly significant.

The general purpose of this investigation, therefore, was to determine the effect of cropping upon the physical and chemical properties of South Dakota soils.

In order to determine the nature and extent of the changes which have taken place in the soil, certain chemical and physical studies of the soils were undertaken. Samples of virgin soils and of cultivated soils were selected from the principal soil types in South Dakota. A comparison of the chemical and physical properties of virgin and of cultivated soils could thus be made in order to determine the possible physical and chemical changes resulting from cultivation.

The specific objects of this investigation may be summarized under the following heads:

I. Physical studies

1. The effect of cultivation upon the size and stability of the soil aggregates or granules.
2. The effect of cultivation upon the mechanical composition of the soil as determined by mechanical analysis.
3. The relationship of the organic matter content of the soil to the size and stability of soil aggregates.

II. Chemical studies

1. The determination of the loss of nitrogen and phosphorus from the soil due to cropping.
2. A study of the organic matter and organic matter-nitrogen ratios of virgin and cultivated soils.
3. A study of the base exchange properties of virgin and cultivated soils.
4. The determination of the soil reaction of virgin and cultivated soils as measured in terms of PH.

PLAN OF INVESTIGATION

Collection of Materials

An attempt was made to obtain samples of the most important soils series in the state. The following series were selected: Barnes, Bearden, Moody, Williams, Orman, and Pierre. Orman and Pierre soils are found west of the

Missouri River in the region of the Chestnuterths. Williams soils lie at the transition zone between Chestnuterths and Chernozems. Barnes, Bearden and Moody soils are all Chernozems.

At this point the differentiation between virgin and cultivated soil is in order. In this study virgin soils are considered any soil which has not been broken with a plow or otherwise cultivated. Its vegetative covering may have been removed for hay or pastured. Cultivated soil is any soil which has been plowed or otherwise broken up and subsequently cropped. Eastern South Dakota soils selected for this study have been cropped for a period of 50 to 55 years.

In gathering samples an attempt was made to select virgin soils from fields which had not been pastured or cut for hay. In all cases, however, this was not possible. When the virgin field was located a sample was taken from a cultivated field lying immediately adjacent. The sample was then taken by carefully cutting a cube of soil to what appeared to be the normal depth of cultivation in that field. A sample was then taken in the same manner to the same depth from the virgin field. In order to eliminate any soil variation due to topography or factors other than cultivation, the samples of virgin and cultivated soils were selected from areas which were in close proximity.

Table 1 gives a description of samples. Samples labeled with the same letter are pairs, the virgin soil of the pair being specified by "sub 1" and the cultivated soil of the pair by "sub 2."

Table 1

DESCRIPTION AND LOCATION OF SAMPLES

Sam- ple	Virgin or Cultivated	Type	Dominant Vegetation	Location
A ₁	Virgin	Pierre clay	<u>Buchloe dactyloides</u> <u>Bouteloua gracilis</u> <u>Opuntia fragilis</u>	Exp. Farm, Cottonwood, S.D.
A ₂	Cultivated	Pierre clay	None	Same as A ₁
B ₁	Virgin	Orman clay	<u>Buchloe dactyloides</u> <u>Bouteloua gracilis</u> <u>Agropyron cristatum</u>	Exp. Farm Cottonwood, S.D.
B ₂	Cultivated	Orman clay	None	Same as B ₁
C ₁	Virgin	Williams clay	<u>Bouteloua gracilis</u> <u>Solsola Kali L.Var.</u> <u>Tenuifolia, C.F.W.</u> <u>Mex Var. Pestifer</u> <u>A. Nels.</u>	Cemetery S.E. of Highmore, S. D.
C ₁	Cultivated	Williams clay loam	None	Same as C ₁
D ₁	Virgin	Barnes sandy loam	<u>Agropyron pauci- florum.</u> <u>Solsola Kali L.Var.</u> <u>Tenuifolia C.F.W.</u> <u>Mex Var. Pestifer</u> <u>A. Nels</u>	Virgin pasture N.W. of Huron, S. D.
D ₂	Cultivated	Barnes sandy	None	Same as D ₁
E ₁	Virgin	Bearden silt loam	<u>A. pauciflorum</u>	Cemetery S.E. of Brookings, S.D.
E ₂	Cultivated	Bearden clay	None	Same as E ₁
F ₁	Virgin	Barnes sandy loam	<u>A. pauciflorum</u>	Cemetery N.W. of Brookings, S.D.
F ₂	Cultivated	Barnes sandy loam	None	Same as F ₁
G ₁	Virgin	Moody silty clay loam	<u>A. pauciflorum</u>	Virgin pasture S. of Brookings, S. Dak.
G ₂	Cultivated	Moody silty clay loam	None	Same as G ₁
H ₁	Virgin	Moody clay loam	<u>A. pauciflorum</u>	Cemetery N.W. of Flandreau, S.D.
H ₂	Cultivated	Moody clay	None	Same as H ₁

DESCRIPTION OF SOIL SERIES

Barnes:

The Barnes series is a glacial soil. The surface of this soil ranges from black to brown in color to a depth of about four to ten inches, resting on a brown soil to a depth of about 18 to 30 inches. Below this a characteristic grayish yellow layer is found. Pebbles and stones are found throughout the soil profile and large boulders may be found even at the surface. The surface soils do not effervesce with hydrochloric acid as a rule, and the lime in the deeper subsoil may be uniformly distributed through the material or may be collected into masses or concretions. (23)

Bearden:

The surface soil of the Bearden series is very dark grayish-brown friable, and finely granular. The surface soil grades into a brown friable subsurface soil which may be heavier than the surface soil. At about two feet in depth the granules disappear, and at from 18 to 22 inches the color changes to yellow. Between depths of about 39 to 48 inches bedded sand and gravel are present, above which a layer of highly calcareous structureless material containing numerous lime concretions occurs. This layer is two to six inches thick. Some of the soils do not have the gravelly substratum and in this case no decided zone of lime accumulation occurs within three feet of the surface. (24).

Moody:

The surface covering of the Moody series is dark grayish-brown to about one inch. Grass roots are abundant in uncultivated soil. Underlying the surface layer is a dark

grayish-brown soil to a depth of from 6 to 12 inches which is turn underlain by a dark grayish-brown or almost black layer (the darkest layer of horizon A) to a depth of from 10 to 18 inches. The undisturbed material is slightly firmer than that of other layers and upon being broken up it becomes lighter in color. The fourth layer is non-calcareous and as a rule slightly heavier than the layers above, ranging in color from dark grayish-brown in the upper part to grayish-brown in the lower. Below a depth of from 30 to 40 inches is the zone of lime accumulation where the material is grayish-yellow in color and structureless. The Moody series is derived from the weathering of loess overlying glacial material at an average depth of about five feet in thickness. It rests on brown and gray mottled clay glacial till, and has a columnar structure. (24).

Williams:

The surface of Williams soil to a depth of about 7 to 12 inches consists of loose, friable, very dark grayish-brown or nearly black soil of finely granular structure. The surface soil is underlain to a depth varying from 16 to 20 inches by dark-brown or brown heavy textured soil containing glacial pebbles. The next layer, which reaches a depth of about 30 inches, is friable or slightly compact, yellow or grayish-yellow in color. It is highly calcareous, containing an abundance of lime in streaks and concretions. Some specks of orange or reddish brown may be present in places. Below this is the pale yellow or yellow friable glacial till from which the soil was derived. (16).

Orman:

DISCUSSION OF RESULTS

The surface soils of Orman series are brown and the subsoils brown or yellowish brown and both have the slaty tinge characteristic of the Pierre series. The series occurs on terraces within or bordering Pierre shale exposures and the material consists mainly of sediments from Pierre soils. The soils are sticky when wet and bake hard when dry. The topography is level and natural drainage is poor on account of the impervious nature of the subsoil.

Pierre:

The soils of the Pierre series are light brown, brown, or olive brown in color. They are compact and refractory when dry and very sticky when wet. The subsoils are brown to olive brown or slaty colored and heavy in texture. They grade into a substratum of partially weathered shale, usually below six feet. The topography is characteristic, consisting of rounded hills and ridges. Surface drainage is usually good. This series is of residual origin, being derived from Pierre shale by weathering.

about that, in the undisturbed soil which is producing a heavy growth of perennial plants, especially the grasses, interfaces are created on the surfaces of the soil granules which differ from the interior of these particles. These interfaces are in many cases largely the result of the action of soil animals and the plants themselves which feed upon their roots. The granules are often so dense that possible organisms probably thrive largely on the surface and not within the granules. If the soil is

DISCUSSION OF RESULTS

The samples had been taken carefully so as to avoid any mechanical change in structure. A representative portion of each sample was removed, air dried, finely ground, and set aside in capped bottles for use in the chemical analysis.

PHYSICAL ANALYSIS

The structural analysis was made upon the unground samples as quickly as possible after they were air dry. Several methods of determining structure have been suggested by various authors. Buehrer (9) suggests the use of the movement of gases through the soil as a measure of soil structure. Rhoades (18) suggests the use of the Kipecky type of elutriator for the separation of different sizes of soil granules. Bayer (3) suggests soil porosity as a measure of aggregation. Yoder (27) and Bouyoucos (5) offer methods based on the water stability of soil particles.

Water stable aggregates are defined and accounted for by Bradfield (8) in his statement that, "In the undisturbed soil which is producing a heavy growth of perennial plants, especially the grasses, interfaces are created on the surface of the soil granules which differ from the interior of these particles. These interfaces are in many cases largely the products or by-products of the plant roots and the micro-organisms which feed upon these roots. The granules are often so dense that aerobic organisms probably thrive largely on the surface and not within the granule. If the soil is

unmolested for a few years these organisms build up an interface that is fairly stable, our so-called 'water resistant' aggregates."

In this work the water stable aggregates were used as a criterion of structure. They were determined in two ways, one method being that of Bouyoucos (4), the other method being similar to that suggested by Yoder (27).

In the Bouyoucos method fifty grams of soil (air dry basis) was placed in a bag of two millimeter window screen and the bag was suspended in a 1000 cc hydrometer cylinder filled with water to the proper mark. The bag was pushed up and down slowly in the water to remove slaked particles. After the soil was completely slaked, five cubic centimeters of normal potassium hydroxide was added and the soil was then shaken six times by placing one hand on the mouth of the cylinder and turning the latter completely upside down and back three times. Hydrometer readings were taken at ten, thirty, and sixty seconds, and at fifteen minutes. The weight of the soil within the various size classes was then determined by referring to the following table:

Time of sedimentation	Largest diameter of particles in suspension
10 sec.	.1914 mm.
30 "	.11009 mm.
60 "	.07785 mm.
15 min.	.00259 mm.

In the Yoder method fifty grams of air dry soil was allowed to slake through a graded nest of screens immersed in a constant volume of water (four liters). The nests of screens were mechanically raised and lowered at a slow rate

to facilitate slaking and separation of the various sized particles. The screens and soil were then dried and the soil was brushed out and weighed in the several size classes.

Structure stability has been measured by Bouyoucos by three methods. In one method he uses the time of slaking and structure of the slaked material as a measure (4). In another method he determines structure stability of the existing aggregates by determining the moisture equivalent of water treated soil and soil treated with potassium chloride and comparing them (7). His third method (7) was used in this study. Two fifty cubic centimeter cylinders were filled to the forty cubic centimeter mark with distilled water and with normal potassium chloride solution respectively. Fifteen grams of air dry soil was added to each cylinder. The soil was first gently loosened by gradually working an iron rod to the bottom of the cylinder and then it was stirred by moving the rod in a forward and backward position twenty times. This stirring was repeated three times at half hour intervals, and the soils were then allowed to stand for twenty-four hours, at which time their settling volume was read. The difference in the settled volume with water and with potassium chloride solution was used as a measure of stability. Stable soils show little difference.

The mechanical analysis was done by the Bouyoucos (6) method.

Aggregate Analysis:


The cause of wind erosion in the Great Plains during recent years has been the subject of much speculation.


Breakdown of structure due to cultivation has been suggested as a possible explanation. One theory which has been advanced is that the decrease in organic matter due to cultivation, together with the mechanical manipulation of the soil by tillage, has caused breakdown of structure. Bradfield (8) states that although they do not ensure favorable structure, calcium and organic matter do aid in the development of such structure. Work to be discussed later indicates a decrease in organic matter through cultivation. If organic matter is largely responsible for the granular nature of the soil, it would seem plausible that a reduction of the organic matter of the soil resulting from cultivation would likewise bring about structural changes in the soil, producing a finer state of aggregation or partial degranulation. The findings of this experiment do not correspond entirely with the above supposition.

In the process of sampling it was noted that in the soils studied the virgin soils showed a more coarsely granular structure and were more friable than the cultivated soils. The coarsely granular structure noted in the virgin soil is no doubt mechanically unstable because as soon as the soil is cultivated the larger aggregates would break down to smaller aggregates such as exist in comparable cultivated soils. It might be presumed that the large visible aggregates, which are relatively unstable, in virgin soils consist of several smaller aggregates.

The methods used for determining the size of water stable aggregates are based on the assumption that the soil

Figure 2.





slakes into its ultimate state of aggregation. It would seem plausible that soils in their natural cultivated state when exposed to rain and mechanical agitation by tillage, could be broken down to the same ultimate state as in the laboratory. It is doubtful if soils are ever completely broken down in their natural state to their ultimate aggregation.

In the soil structure studies it was found that the most suitable method was the determination of water stable aggregates of various sizes since it seemed that this was the only uniform measurement which could be made.

Water stable aggregates were determined by the Bouyoucos and Yoder methods discussed previously. Tables 2 and 3 show detailed results of the two methods. Figures 2 and 3 show the comparison of the averages of the several aggregate analysis on the virgin and cultivated soils.

Because there is no universally accepted method for the determination of the size of soil aggregates, the accurate measurement of soil structure becomes a difficult problem. However, the methods available are capable of determining significant differences of structure between various soils.

The Yoder method measures larger aggregates than the Bouyoucos method, and this must be taken into consideration when figures 2 and 3 are examined. A larger per cent of the largest granules were found in the virgin soil than in the cultivated. Some difference is found in the amounts of smaller aggregates, the cultivated soil showing slightly higher percentages than the virgin soil.

Results of the structural analysis by the Bouyoucos method shown in figure 3 show a slight difference in the virgin and cultivated soil, the virgin soil having the larger percentage of larger aggregates and smaller percentage of small aggregates, which agrees with the results of the Yoder method.

Both sets of results show some change in structure, but the change is small, so small in fact that it appears the relation of structure to wind erosion may have been over-emphasized.

Table 2

RESULTS OF THE BOUYOUCOS METHOD OF AGGREGATE ANALYSIS

Sample	Over .1914 mm. %	.1914- .11009 mm. %	.11009- .007785 mm. %	.007785 .00259 mm. %	.00259 .0000 mm. %
Virgin Soil Type					
A ₁ Pierre Clay	82.32	5.00	1.00	8.28	3.40
B ₁ Orman Clay	70.64	10.00	5.00	12.36	2.00
C ₁ Williams Clay	71.60	9.00	5.00	11.84	2.56
D ₁ Barnes Sandy loam	74.52	14.00	2.00	7.64	1.84
E ₁ Bearden Silt loam	52.00	11.00	13.00	21.88	2.12
F ₁ Barnes Sandy loam	76.16	7.00	8.00	4.28	4.56
G ₁ Moody Silty Clay loam	67.12	8.00	6.56	14.92	3.40
H ₁ Moody Clay loam	<u>62.28</u>	<u>12.50</u>	<u>7.52</u>	<u>14.14</u>	<u>3.56</u>
Total	556.64	76.50	48.08	95.34	23.44
Av.	69.58	9.56	6.01	11.92	2.93

Table 2 (continued)

RESULTS OF THE BOUYOUCOS METHOD OF AGGREGATE ANALYSIS (continued)

Sample	Size of Aggregates				
	Over .1914 mm. %	.1914- .11009 mm. %	.11009- .007785 mm. %	.007785- .00259 mm. %	.00259- .0000 mm. %
Cultivated					
Soil Type					
A ₂ Pierre Clay	58.32	10.00	12.00	16.00	3.68
B ₂ Orman Clay	50.32	12.00	12.00	21.68	4.00
C ₂ Williams Clay loam	67.60	8.00	4.00	15.28	5.12
D ₂ Barnes Sandy loam	72.72	4.00	8.00	9.64	5.64
E ₂ Bearden Clay loam	77.32	10.00	3.00	7.28	2.40
F ₂ Barnes Sandy loam	76.38	5.00	4.00	9.78	4.84
G ₂ Moody Silty Clay loam	62.12	8.00	6.00	19.20	4.68
H ₂ Moody Clay loam	<u>51.88</u>	<u>11.50</u>	<u>7.50</u>	<u>23.92</u>	<u>5.20</u>
Total	516.66	68.50	56.50	122.78	35.56
Av.	64.58	8.56	7.06	15.35	4.45

Table 3

RESULTS OF THE YODER METHOD OF AGGREGATE ANALYSIS

Sample	Size of Aggregates in mm.					
	.4200-	.2500-	.1770-	.1490-	.0740	.0740-
	.8400	.4200	.2500	.1770	.1490	.0000
	%	%	%	%	%	%
Virgin Soil Type						
A ₁ Pierre Clay	43.02	22.84	5.83	1.79	15.18	11.34
B ₁ Orman Clay	42.79	21.76	8.22	3.80	14.68	8.75
C ₁ Williams Clay	40.64	31.13	3.08	0.63	11.97	12.55
D ₁ Barnes Sandy loam	42.98	35.32	2.97	0.65	9.42	8.66
E ₁ Bearden Silt loam	44.71	16.08	12.12	2.01	8.06	17.02
F ₁ Barnes Sandy loam	28.93	25.34	8.30	2.55	15.42	19.46
G ₁ Moody Silty Clay loam	36.89	21.80	5.96	0.84	19.21	15.30
H ₁ Moody Clay loam	<u>47.41</u>	<u>14.35</u>	<u>8.35</u>	<u>1.77</u>	<u>9.85</u>	<u>18.27</u>
Total	327.37	188.62	54.83	14.04	103.79	111.35
Av.	40.92	23.58	6.85	1.76	12.97	13.92

Table 3 (continued)

RESULTS OF THE YODER METHOD OF AGGREGATE ANALYSIS (continued)

Sample	Size of Aggregates in mm.					
	.4200- .8400 %	.2500- .4200 %	.1770- .2500 %	.1490- .1770 %	.0740- .1490 %	.0740 .0000 %
Cultivated						
Soil Type						
A ₂ Pierre Clay	31.79	43.31	3.45	0.65	8.14	12.16
B ₂ Ormen Clay	4.59	19.36	28.41	11.64	15.90	20.10
C ₂ Williams Clay loam	26.37	32.46	8.22	1.77	14.02	17.16
D ₂ Barnes Sandy loam	33.19	33.13	3.35	1.30	14.77	14.26
E ₂ Bearden Clay loam	38.21	18.59	13.09	0.95	10.95	18.21
F ₂ Barnes Sandy loam	38.62	29.04	7.68	1.54	8.10	15.02
G ₂ Moody Silty C Clay loam	27.74	50.16	2.04	0.85	7.35	11.86
H ₂ Moody Clay loam	<u>24.33</u>	<u>13.15</u>	<u>24.63</u>	<u>2.24</u>	<u>12.67</u>	<u>22.98</u>
Total	224.84	239.20	90.87	20.94	91.90	132.25
Av.	28.10	29.90	11.36	2.61	11.50	16.53

Structural Stability

Table 4 shows the results obtained by the use of the Bouyoucos method for determining structural stability of the soil aggregates in water. Stable structure is shown by a small difference in the volume of the soil in water as compared to its volume in potassium chloride solution. The potassium chloride prevents the breakdown of structural aggregates by keeping the colloidal material in its flocculated state. If the soil aggregates are not stable in water, when placed in water they will be dispersed. Dispersion of the aggregates

will result in the formation of smaller particles, and as the size of particles decreases the volume of the soil increases. Those soils in the potassium chloride solution will retain the volume of the soil in which the aggregates have not been dispersed, and the difference in volume of the differently treated soils can therefore be used as an index of water stability.

Table 4

RESULTS OF THE STUDY OF STABILITY
OF AGGREGATES ON VIRGIN AND CULTIVATED SOIL
BY THE BOUYOUCOS METHOD

Sample	Soil Type	Volume in Water cc.	Volume in KCL Soln. cc.	Difference in Volume cc.
A ₁	Pierre Clay	18.5	17.0	1.5
A ₂	Pierre Clay	16.0	16.0	0.0
B ₁	Orman Clay	17.3	16.8	0.5
B ₂	Orman Clay	17.0	16.5	0.5
C ₁	Williams Clay	18.5	18.0	0.5
C ₂	Williams Clay loam	16.5	17.0	-0.5
D ₁	Barnes Sandy loam	16.0	16.0	0.0
D ₂	Barnes Sandy loam	14.0	14.0	0.0
E ₁	Bearden Silt loam	20.3	21.8	-1.5
E ₂	Bearden Clay loam	17.8	17.3	0.5
F ₁	Barnes Sandy loam	19.5	19.5	0.0
F ₂	Barnes Sandy loam	15.5	16.0	-0.5
G ₁	Moody Silty Clay loam	17.5	18.0	-0.5
G ₂	Moody Silty Clay loam	17.3	17.3	0.0
H ₁	Moody Clay loam	20.5	20.5	0.0
H ₂	Moody Clay loam	17.3	17.3	0.0

In this work a soil showing a difference in volume of less than 2 cubic centimeters was considered stable. From table 4 it may be concluded that there is no difference in the water stability of structure between cultivated and virgin soils. The soil used in the structural stability studies belong in the Pedocal group and consequently they have colloidal complexes saturated with calcium. Soils whose

colloidal complexes are saturated with calcium should have a stable structure and resist any mechanical breakdown when placed in water. The results obtained in the structural stability studies agree with the above facts since all the soils possessed a water stable structure. Cultivation had little or no effect on the water stability of the granules. It may have been assumed that the loss of organic matter would have some effect on the stability of the granules because of the decrease in the amount of humus and consequent decrease in the organic colloids which together with inorganic colloids are considered responsible for binding the soil particles in granules. It may be presumed that the humus or organic colloids remaining in the soil are sufficient to stabilize the soil granules against the dispersing or degrenulating effects of water such as the soils are subjected to in the Bouyoucos method for determining water stability.

In the Bouyoucos method for determining state of aggregation of the soils, fifty grams of soil was allowed to slake through a 2 mm. mesh screen by slowly lowering and raising the soil and screen in 1000 cc of water. The time required for the soil to pass through the screen is considered the time required to slake. The following results were obtained.

Virgin Soil		Cultivated Soil	
Sample	Time to Slake	Sample	Time to Slake
A ₁ -----	10 min.-----	A ₂ -----	1 min.-----
B ₁ -----	3 min.-----	B ₂ -----	1 min.-----
C ₁ -----	10 min.-----	C ₂ -----	12 min.-----
D ₁ -----	19 min.-----	D ₂ -----	5 min.-----
E ₁ -----	16 min.-----	E ₂ -----	14 min.-----
F ₁ -----	13 min.-----	F ₂ -----	5 min.-----
G ₁ -----	12 min.-----	G ₂ -----	9 min.-----
H ₁ -----	28 min.-----	H ₂ -----	5 min.-----

The results of the slaking experiment show that the virgin soils require as an average 13.9 minutes to slake and the cultivated soils require 6.5 minutes. This difference may be attributed in part to the presence of roots in the virgin soil which tend to hold the soil together.

Mechanical Analysis

Daniel (1) made several mechanical analyses to compare virgin and cropped with drift soil in the Southern High Plains. He found very little difference in the sand, silt, and clay content of virgin and cultivated soil.

Table 5 shows the results of the mechanical analysis of the soils used in the work presented in this paper.

Table 5

RESULTS OF THE MECHANICAL ANALYSIS OF VIRGIN AND CULTIVATED SOUTH DAKOTA SOILS

Sample	% Sand	% Silt	% Clay	Class	<u>Sand & Silt</u> Clay
A ₁	24.98	27.82	47.20	Clay	1.119
A ₂	25.60	30.04	44.36	Clay	1.254
B ₁	18.00	24.72	57.28	Clay	0.746
B ₂	18.52	26.20	55.28	Clay	0.809
C ₁	23.60	44.84	31.56	Clay	2.169
C ₂	34.16	39.56	26.28	Clay loam	2.805
D ₁	57.44	23.36	19.20	Sandy loam	4.208
D ₂	56.96	23.68	19.36	Sandy loam	4.165
E ₁	30.96	50.84	18.20	Silt loam	4.495
E ₂	39.48	38.40	22.12	Clay loam	3.521
F ₁	60.84	28.60	10.56	Sandy loam	8.470
F ₂	64.18	19.10	16.72	Sandy loam	4.981
G ₁	23.88	55.20	20.92	Silty clay loam	3.780
G ₂	22.26	52.62	25.12	Silty clay loam	2.981
H ₁	35.86	42.22	21.92	Clay loam	3.562
H ₂	32.16	40.08	27.76	Clay loam	2.602

From the table it is seen that the cultivated clay soils have less clay and more sand than the virgin clay soils, the silt varying each way. Those soils having a lower clay content have increased clay percentages in the cultivated soils.

Figure 4 shows a comparison of the sand + silt ratios of clay the virgin and cultivated soils. It is possible that there may have been some variation in the original textures of the cultivated and virgin soils of a pair, i.e., before the sod was broken the soil at the location from which the cultivated sample was taken may have varied in texture from the soil at the location where the virgin sample was taken. However, the samples were taken in such close proximity that variations as large as are indicated by the analysis could not be due to this factor alone.

In the soils of coarser texture the increase in clay content is probably due to the fact that percolation of water through the soil has carried the finer materials from the surface downward. Erosion of the surface soil, and cultivation would then result in the mixture of the surface with the sub-surface soils which contain a higher per cent of clay. In the clay soils, because of their impervious nature and because they lie in an area of such low rainfall, removal of clay from the surface by mechanical elevation to the subsurface has not occurred to any great extent. Erosion by wind tends to remove from these clay soils a larger per cent of clay than of silt and sand, causing the increased sand + silt ratio.
clay

Figure 4

A COMPARISON OF SAND + SILT
CLAY
RATIOS OF CULTIVATED AND
VIRGIN SOUTH DAKOTA SOILS.


Legend



Virgin soil



Cultivated soil



CHEMICAL ANALYSIS

Moisture was determined on the ground sample by heating at 110° centigrade for twenty-four hours. Loss on ignition was then determined by heating at a red heat till all organic material was burned off.

Organic matter was determined by the hydrogen peroxide method suggested by Robinson (15).

Humus was determined by extraction with a weak alkali. 10 grams of air dry soil was leached with one per cent hydrochloric acid until free of calcium. The hydrochloric acid was then washed out with water, the soil and the filter paper dried, and when dry the soil and filter paper were placed in a shaker bottle. Two hundred fifty cubic centimeters of four per cent ammonium hydroxide was then added and the soil was shaken for three hours. The soil was then allowed to settle and fifty cubic centimeters of the filtered supernatant liquid was evaporated to dryness, dried in an oven at 110° centigrade, and weighed. It was then burned at a red heat till all organic matter was oxidized, cooled, and weighed. The difference in the two weights represents the humus.

Nitrogen was determined by the Kjeldahl method. PH was determined with a glass electrode outfit, the method being that of Deen and Walker (12). Phosphorus was determined by the A. O. A. C. method (1). Base exchange capacity was found by the Parker method as described by Walker, Brown, and Young (14).

Loss on Ignition

Loss on ignition was determined on the soils as an indication of organic matter only.

Table 6

LOSS ON IGNITION OF THE VIRGIN AND CULTIVATED SOILS

Sample	Soil Type	Per cent loss on ignition	Loss through Cultivation*	Per cent gain or loss, based on the virgin soil*
A ₁	Pierre Clay	7.795		
A ₂	Pierre Clay	6.088	-1.707	-21.90
B ₁	Orman Clay	6.742		
B ₂	Orman Clay	6.546	-0.196	- 2.91
C ₁	Williams Clay	8.859		
C ₂	Williams Clay loam	6.371	-1.988	22.44
D ₁	Barnes Sandy loam	6.657		
D ₂	Barnes Sandy loam	5.492	-1.165	-17.50
E ₁	Bearden Silt loam	11.829		
E ₂	Bearden Clay loam	8.514	-3.315	-28.02
F ₁	Barnes Sandy loam	10.772		
F ₂	Barnes Sandy loam	4.987	-5.785	-53.70
G ₁	Moody Silty Clay loam	10.194		
G ₂	Moody Silty Clay loam	8.977	-1.217	-11.94
H ₁	Moody Clay loam	11.182		
H ₂	Moody Clay loam	8.608	-2.574	-23.02

*Gains, in tables similar to this, are represented by plus, and losses by minus.

An examination of the figures shows that in all cases the loss on ignition is lower in the cultivated than in the virgin soil.

Organic Matter

The importance of the amount and quality of soil organic matter in relation to crop growth has long been recognized. The functions of organic matter in the soil may be classed as physical, chemical, biological, and biochemical.

The physical effects of organic matter upon soil are extremely complex. From the physical standpoint organic matter affects the properties of soil in the following respects: weight, cohesion, structure, absorption, porosity, color, temperature, and tilth. Any change in the amounts of organic matter in the soil may affect one or all of the above physical properties. The present study has attempted to correlate the loss of organic matter with structural changes in the soil.

The chemical, biological, and biochemical effects of organic matter are of considerable importance from the standpoint of soil fertility. Chemically organic matter functions in three important ways, namely: (1) organic matter constitutes the most valuable source of soil nitrogen; (2) through biochemical changes or decomposition of organic matter mineral plant food is made available; (3) organic matter functions in base exchange and other soil reactions. It is obvious that losses of soil organic matter will have a decided influence upon soil fertility and consequently effect crop production.

approximately 42 per cent of the amount contained in the virgin soil. These figures indicate that during the relatively short period in which South Dakota soils have been cropped almost one-half of the original organic matter has been depleted.

Table 7

THE ORGANIC MATTER CONTENT
OF VIRGIN AND CULTIVATED SOILS

Sample Soil Type		Per cent organic matter	Gain or loss through cultivation	Per cent gain or loss based on the virgin soil content
A ₁	Pierre Clay	3.706		
A ₂	Pierre Clay	2.386	-1.320	-35.62
B ₁	Orman Clay	2.478		
B ₂	Orman Clay	1.399	-1.079	-43.54
C ₁	Williams Clay	5.978		
C ₂	Williams Clay loam	3.389	-2.089	-34.94
D ₁	Barnes Sandy loam	4.035		
D ₂	Barnes Sandy loam	3.202	-0.833	-20.64
E ₁	Bearden Silt loam	9.060		
E ₂	Bearden Clay loam	4.707	-4.353	-48.05
F ₁	Barnes Sandy loam	9.573		
F ₂	Barnes Sandy loam	3.363	-6.210	-64.87
G ₁	Moody Silty Clay loam	6.455		
G ₂	Moody Silty Clay loam	4.912	-1.543	-23.90
H ₁	Moody Clay loam	9.472		
H ₂	Moody clay loam	5.404	-4.068	-42.95

A study of table 7 reveals that losses of organic matter resulted in all eight of the cultivated soils studied. The loss of organic matter was found to be greater than the loss of any other soil constituent determined in this study. The average loss of organic matter through cultivation was approximately 42 per cent of the amount contained in the virgin soil. Thus these figures indicate that during the relatively short period in which South Dakota soils have been cropped almost one-half of the original organic matter has been depleted.

From the structural studies it was concluded that cultivation caused little change in the soil structure. Losses in organic matter, however, have been shown to be large. These two facts are not in agreement with the ordinary conception of the relationship which exists between organic matter and structure.

The determination of the effect of the quantity of organic matter, in the soil upon the soil structure involves certain difficulties. It is entirely possible that a certain definite quantity or percentage of organic matter brings about the production of the highest state of granulation as measured by the size or stability of the granules. From a study of the results obtained it might be concluded that the reduction in organic matter due to cultivation has not been large enough to have a decided influence upon the size and stability of the granules, but it may be the cause of decreased resistance to water slaking found in the Bouyoucos method for determining soil structure.

Humus

Humus, as determined in this study, may be defined as that portion of the organic matter which is soluble in a weak base. Table 8 gives the results of the humus determination.

The results of the nitrogen determinations on the

cultivated and virgin soils are shown in table 9.

Table 8

THE HUMUS CONTENT OF THE VARIOUS
VIRGIN AND CULTIVATED SOILS

Sample	Soil Type	% Humus	Loss through cultivation	Per cent loss, based on the humus content of the virgin soil
C ₁	Williams Clay	5.110		
C ₂	Williams clay loam	2.645	2.465	48.24
D ₁	Barnes sandy loam	2.870		
D ₂	Barnes sandy loam	2.575	0.295	10.28
E ₁	Bearden silt loam	5.285		
E ₂	Bearden clay loam	3.870	1.415	26.77
F ₁	Barnes sandy loam	4.255		
F ₂	Barnes sandy loam	2.333	1.922	45.12
G ₁	Moody silty clay loam	3.770		
G ₂	Moody silty clay loam	3.515	0.255	6.76
H ₁	Moody clay loam	3.765		
H ₂	Moody clay loam	2.810	0.955	25.37

Humus on samples A and B was not determined because of the effect upon the results of the high colloidal content of the soils.

In the six soils shown in table 8 losses in humus were found in all cases, the average loss being 29.163 per cent, when the humus content of the virgin soil is used as the basis for computing the losses.

Nitrogen

The results of the nitrogen determinations on the cultivated and virgin soils are shown in table 9.

Percent Loss, Based on the Virgin Soil




Table 9

THE NITROGEN CONTENT OF THE VIRGIN
AND CULTIVATED SOILS

Sample	Soil Type	Per cent Nitrogen	Gain or loss through cultivation	Per cent gain or loss based on the virgin soil
A ₁	Pierre Clay	0.1877		
A ₂	Pierre Clay	0.1267	-0.0610	-32.50
B ₁	Orman clay	0.1200		
B ₂	Orman clay	0.1106	-0.0094	- 7.83
C ₁	Williams clay	0.2624		
C ₂	Williams clay loam	0.1798	-0.0826	-31.48
D ₁	Barnes sandy loam	0.1869		
D ₂	Barnes sandy loam	0.1494	-0.0375	-20.06
E ₁	Bearden silt loam	0.4090		
E ₂	Bearden clay loam	0.2690	-0.1400	-34.23
F ₁	Barnes sandy loam	0.3795		
F ₂	Barnes sandy loam	0.1555	-0.2240	-59.03
G ₁	Moody silty clay loam	0.3251		
G ₂	Moody silty clay loam	0.2760	-0.0491	-15.10
H ₁	Moody clay loam	0.4059		
H ₂	Moody clay loam	0.2739	-0.1320	-32.52

A loss in nitrogen occurred in every case as the result of cultivation. The average loss for all soils, using virgin soil as the base, is about 29 per cent, indicating that unless cultural practices are changed South Dakota farmers will within a relatively short period have confronting them the additional problem of nitrogen shortage. From figure 5 it

will be seen that losses in nitrogen and loss on ignition are similar, the percentage losses in nitrogen being slightly larger in all cases. Losses in humus and organic matter also correspond closely with losses in nitrogen. A study of table 10 reveals that the organic matter-nitrogen ratio (total organic matter divided by the nitrogen) is wider in the virgin soil than in the cultivated soils. This would indicate that soils lose organic matter at a more rapid rate than nitrogen.

Table 10

A COMPARISON OF THE ORGANIC MATTER, NITROGEN, AND HUMUS CONTENTS OF THE VARIOUS SOILS, AND THE ORGANIC MATTER-NITROGEN RATIOS OF THE VIRGIN AND CULTIVATED SOILS

Sample	Soil Type	Per cent organic matter	Per cent Nitrogen	Organic matter divided by nitrogen	Per cent Humus
A ₁	Pierre Clay	3.706	.1877	19.744	---
A ₂	Pierre Clay	2.386	.1267	18.832	---
B ₁	Orman clay	1.870	.1200	15.583	---
B ₂	Orman clay	1.102	.1106	9.964	---
C ₁	Williams clay	5.978	.2624	22.782	5.110
C ₂	Williams clay loam	3.889	.1798	21.630	2.645
D ₁	Barnes sandy loam	4.035	.1869	21.589	2.870
D ₂	Barnes sandy loam	3.202	.1494	21.432	2.575
E ₁	Bearden silt loam	9.060	.4090	22.152	5.285
E ₂	Bearden clay loam	4.707	.2690	17.498	3.870
F ₁	Barnes sandy loam	9.573	.3795	25.225	4.255
F ₂	Barnes sandy loam	3.363	.1555	21.627	2.333
G ₁	Moody silty clay loam	6.455	.3251	19.855	3.770
G ₂	Moody silty clay loam	4.912	.2760	17.797	3.515
H ₁	Moody clay loam	9.472	.4059	23.336	3.765
H ₂	Moody clay loam	5.404	.2739	19.730	2.810

Phosphorus

A comparison of virgin and cultivated soils with regard to phosphorus content is shown in table 11. Three soils show gains and five show losses in phosphorus as the result of cultivation. DeTurk (13) states that crop residues return phosphorus mainly to the surface soil while plant roots remove it from the surface and also from greater depths. Following partial or total crop failures of Western South Dakota, the unharvested crops have frequently been left upon the fields. All of the phosphorus which has been removed from the surface as well as that removed from greater depths by crops, has thus been concentrated in the surface soil. With repeated crop failures the phosphorus content of the surface soil is actually increased. Increases in sample A (Pierre Clay) and B (Orman Clay) may be accounted for in this manner.

Table 11

COMPARISON OF THE PHOSPHORUS CONTENT OF VIRGIN AND CULTIVATED SOILS

Sample	Soil Type	Per cent phosphorus	Gain or loss through cultivation	Per cent gain or loss, based on the virgin soil
A ₁	Pierre Clay	0.0832		
A ₂	Pierre Clay	0.0942	+0.0110	+13.22
B ₁	Orman Clay	0.1028		
B ₂	Orman Clay	0.1053	+0.0025	+ 2.43
C ₁	Williams Clay	0.0771		
C ₂	Williams Clay loam	0.0711	-0.0060	- 7.78
D ₁	Barnes Sandy loam	0.0746		
D ₂	Barnes Sandy loam	0.0783	+0.0037	+ 4.95

Table 11 (continued)

COMPARISON OF THE PHOSPHORUS CONTENT
OF VIRGIN AND CULTIVATED SOILS (continued)

Sample	Soil Type	Per cent phosphorus	Gain or loss through cultivation	Per cent gain or loss, based on the virgin soil
E ₁	Bearden Silt loam	0.1098		
E ₂	Bearden Clay loam	0.0910	-0.0138	-17.12
F ₁	Barnes Sandy loam	0.0793		
F ₂	Barnes Sandy loam	0.493	-0.0300	-37.83
G ₁	Moody Silty Clay loam	0.1126		
G ₂	Moody Silty Clay loam	0.1093	-0.0033	- 2.93
H ₁	Moody Clay loam	0.1181		
H ₂	Moody Clay loam	0.0970	-0.0211	-17.87

In only one case, sample D, was the phosphorus content of the cultivated soil higher than the virgin soil in the Chernozems (dark-colored soil) of eastern South Dakota. In all other cases the phosphorus content of the soil has been increased as the result of cultivation. The average decrease in phosphorus for the eastern South Dakota soils is 13.1 per cent. The largest loss of phosphorus was found in sample F which is a sandy loam. It is possible that the large decrease is due to the fact that a large part of the phosphorus is contained in the organic fraction of the soil, the mineral fraction having a relatively low content of the element, and oxidation of the organic matter has released large amounts of the phosphorus which have been used by plants or leached out of the surface layer.

Soil Reaction

Soil Reaction of the soils was determined as PH with a glass electrode potentiometer. From table 12 it is evident that in the eastern South Dakota soils the PH has been decreased (acidity increased) as result of cultivation in all cases except one (sample D) where it remained constant. In the soils from western and central South Dakota small increases are noted in samples B and C. In samples A and B the calcium carbonate occurs fairly near the surface in large enough amounts to effervesce with hydrochloric acid. In sample C the zone of calcium carbonate occurs fairly near the surface. Thus, with plowing, an increase in PH might be expected due to mixing of the horizons. The high root content of the virgin soil may act also in giving it a lower PH than the cultivated soil. In eastern South Dakota the rainfall is great enough so that the calcium carbonate layer is considerably deeper beneath the surface of the soil. The mixing of the horizons by cultivation to the ordinary depths would not cause decreased acidity.

Table 12

A COMPARISON OF THE PH OF VIRGIN AND CULTIVATED SOUTH DAKOTA SOILS

	Sample							
	A	B	C	D	E	F	G	H
Virgin soil-----	7.50	8.20	6.40	6.60	6.40	7.10	6.30	7.00
Cultivated soil-----	7.25	8.50	6.70	6.60	6.30	6.85	5.90	6.50


Figure 6 shows the differences in PH values in the cultivated and virgin soils, and also shows the differences in changes in PH values in western and eastern South Dakota. Sample D is

A COMPARISON OF THE pH VALUES OF VIRGIN AND CULTIVATED SOUTH DAKOTA SOILS

LEGEND

— Virgin soil

--- Cultivated soil



the point of change. Sample D was taken just east of the twenty inch rainfall line.

Base Exchange Capacity

Base exchange capacity of the soil is dependent upon the colloidal content, including both the mineral and organic colloidal fractions. Although not all chemical constituents of the soil take part in base exchange, it is one of the important principles concerned in soil fertility.

The base exchange capacities of the cultivated soils studied are given in table 13. Two of the soils showed increases and six showed decreases as the result of cultivation. Decreases ranged from 6.840% to 43.383% (using virgin soil as the base.) while the two gains were 3.677% and 5.984%. The gains were so small that they cannot be considered highly significant.

Table 13

A COMPARISON OF THE BASE EXCHANGE CAPACITIES OF THE VIRGIN AND CULTIVATED SOILS

Sample	Soil Type	Base exchange capacity M.E. per 200 gm. soil	Gain or loss due to culti- vation	Per cent gain or loss based on the virgin soil
A ₁	Pierre Clay	25.833		
A ₂	Pierre Clay	27.353	+1.520	+5.884
B ₁	Orman Clay	33.451		
B ₂	Orman Clay	31.163	-2.288	-6.840
C ₁	Williams Clay	23.358		
C ₂	Williams Clay loam	18.462	-4.896	-20.961
D ₁	Barnes Sandy loam	16.619		
D ₂	Barnes Sandy loam	15.162	-1.457	-8.767
E ₁	Bearden Silt loam	27.254		
E ₂	Bearden Clay loam	21.513	-5.741	-21.065

Table 13 (continued)

A COMPARISON OF THE BASE EXCHANGE CAPACITIES
OF THE VIRGIN AND CULTIVATED SOILS (continued)


Sample	Soil Type	Base exchange capacity M.F. per 200 gm. soil	Gain or loss due to culti- vation	Per cent gain or loss based on the virgin soil
F ₁	Barnes Sandy loam	24.947		
F ₂	Barnes Sandy loam	12.877	-12.070	-48.383
G ₁	Moody Silty Clay loam	22.572		
G ₂	Moody Silty Clay loam	23.402	+ 0.830	+ 3.677
H ₁	Moody Clay loam	26.155		
H ₂	Moody Clay loam	23.898	+ 2.257	+ 8.629

Figure 7 compares losses in organic matter and base exchange capacity, showing a definite relationship between the two. Large losses in organic matter are accompanied by large losses in base exchange capacity.

Since organic matter possesses high base exchange capacity, any loss of organic matter would reduce the base exchange capacity of the soil.

The correlation between the base exchange capacity and the organic matter content are consistent with the generally accepted fact that organic colloids are dynamic factors in base exchange capacity.

A Comparison of Percent Losses of Base Exchange Capacity and Organic Matter of South Dakota Soils, Due to Cultivation



THE CHEMICAL ANALYSIS OF THE VARIOUS SOILS USED IN THE STUDY

Sam- ple	Soil Type	Mois- ture	Loss on Ignition	Humus	Organ- ic mat- ter	PH	Phos- phorus	Nitro- gen	Base Ex- change Capacity	MF. per 100 gm. soil
		%	%	%	%	%	%	%		
A ₁	Pierre Clay	3.500	7.795		3.706	7.50	.0832	.1877	25.823	
A ₂	Pierre Clay	3.385	6.088		2.386	7.25	.0942	.1267	27.353	
B ₁	Orman Clay	4.295	6.742		2.478	8.20	.1028	.1200	33.451	
B ₂	Orman Clay	3.835	6.546		1.399	8.50	.1053	.1106	31.163	
C ₁	Williams Clay	2.900	8.859	5.110	5.978	6.40	.0771	.2624	23.358	
C ₂	Williams Clay loam	1.895	6.871	2.645	3.889	6.70	.0711	.1798	18.462	
D ₁	Barnes Sandy loam	2.600	6.657	2.870	4.035	6.60	.0746	.1869	16.619	
D ₂	Barnes Sandy loam	2.560	5.492	2.575	3.202	6.60	.0783	.1494	15.162	
F ₁	Bearden Silt loam	2.105	11.829	5.285	9.060	6.40	.1098	.4090	27.254	
F ₂	Bearden Clay loam	1.645	8.514	3.870	4.707	6.30	.0910	.2690	21.513	
F ₁	Barnes Sandy loam	1.860	10.772	4.255	9.573	7.10	.0793	.3795	24.947	
F ₂	Barnes Sandy loam	0.975	4.987	2.333	3.363	6.85	.0493	.1555	12.877	
G ₁	Moody Silty Clay loam	2.240	10.194	3.770	6.455	6.30	.1126	.3251	22.572	
G ₂	Moody Silty Clay loam	2.595	8.977	3.515	4.912	5.90	.1093	.2760	23.402	
H ₁	Moody Clay loam	2.400	11.182	3.765	9.472	7.00	.1181	.4059	26.155	
H ₂	Moody Clay loam	2.380	8.608	2.810	5.404	6.50	.0970	.2739	23.898	


SUMMARY

1. As determined by the Bouyoucos and Yoder methods, the ultimate structure of the cultivated soils was found to be little changed.
There were no highly significant differences in the distribution of the various sizes of the soil aggregates, as measured by the Bouyoucos or Yoder method, in either the virgin or cultivated soils.
Some breakdown of structure was evidenced by the higher per cent of the larger aggregates and the lower per cent of smaller aggregates in the virgin soil, according to the results of both methods.
2. Water stability of aggregates, as measured by the Bouyoucos method, was found to be the same in virgin and in cultivated soils, but cultivated soils showed a tendency toward more rapid slaking in water than virgin soils.
3. Losses in humus and loss on ignition occurred in the cultivated soils in every case.
4. Organic matter is known to aid in the development of good soil structure. An average loss of 42 per cent of the original organic matter resulted from cultivation in the soils which were studied. With such large losses of organic matter changes in structure would be anticipated. Since marked changes in structure were not found, it appears that while organic matter may be necessary in the development and maintenance of favorable structure,

its value diminishes after the soil has acquired sufficient organic matter for the development of maximal structure.

In the virgin soils, the larger amounts of organic matter were probably responsible for the increased time of slaking.

5. The mechanical analyses showed that little change in the texture of the soils has been brought about by cultivation. In three clay soils studied, a slight increase in the sand + silt ratio resulted. All other soils showed slight clay decreases in the sand + silt ratios of the cultivated soils.
clay
6. Cultivation resulted in a decrease in base exchange capacity in all soils studied with the exception of two. Since these two represent relatively small increases as compared with the decreases, it may be that original differences in the soil were responsible for the apparent discrepancy. Decreases in base exchange capacity were found to correspond closely with decreases in organic matter.
7. Losses in nitrogen were found in all cases. The average loss for the eight cultivated soils was found to be 29 per cent of the original nitrogen content. The organic matter ratio was however found to be decreased nitrogen by cultivation, indicating a higher concentration of nitrogen in the organic matter of cultivated soils.
8. The phosphorus content of three soils was increased through cultivation. Decreases resulted in all other cultivated soils studied. In general, the eastern South Dakota soils decreased in phosphorus content as the result of cultivation.



LITERATURE CITED

9. The pH of western South Dakota soils was increased, while that of eastern South Dakota soils was decreased, as the result of cultivation.
10. Figure 8 summarizes the chemical changes of South Dakota soils brought about by cultivation.
1. Reyer, L. E.
"Soil porosity as an index of structure."
Soil Survey Assoc. Trans. 19:71-86 1921
2. Bouyoucos, C. J.
"Texture and structure of soils as influenced by chemical agents."
Jour. Ag. Res. Assoc. 19:793-99 1927
3. Bouyoucos, C. J.
"The ultimate natural structure of the soil."
Soil Science 25:17-37 1929
4. Bouyoucos, C. J.
"A comparison between the pipette method and the hydrometer method for making mechanical analysis of soils."
Soil Science 18:135-43 1924
5. Bouyoucos, C. J.
"Simple and rapid methods for ascertaining the existing structural stability of soil aggregates."
Journal Ag. Res. Assoc. 17:723-7 1925
6. Bradfield, Richard
"The value and limitations of sodium in soil structure."
Soil Survey Assoc. Trans. 17:31-4 1926
7. Buchner, E. P.
"The movement of gases through the soil as a criterion of soil structure."
Agric. Sci. Exp. Sta. Trans. Bull. No. 39 1932
8. Daniel, H. J.
"The physical changes in soils of the Southern High Plains due to cropping and wind erosion and the relation between the sand & silt ratios in these soils."
Jour. Ag. Res. Assoc. 19:573-80 1926

LITERATURE CITED

1. -----
Official and Tentative Methods of Analysis of the
Association of Official Agricultural Chemists.
Second Edition 1924
2. -----
"Summary of observations of field men on cultural
changes in soils."
Am. Soil Survey Assn. Rpt. 12:71-86 1931
3. Bayer, L. D.
"Soil porosity as an index of structure."
Am. Soil Survey Assn. Rpt. 14:83-5 1933
4. Bouyoucos, G. J.
"Texture and structure of soils as influenced by
chemical agents."
Jour. Am. Soc. Agron. 19:788-97 1927
5. Bouyoucos, G. J.
"The ultimate natural structure of the soil."
Soil Science 28:27-37 1929
6. Bouyoucos, G. J.
"A comparison between the pipette method and the
hydrometer method for making mechanical analysis
of soils."
Soil Science 38:335-43 1934
7. Bouyoucos, G. J.
"Simple and rapid methods for ascertaining the
existing structural stability of soil aggregates."
Journal Am. Soc. Agron. 27:222-7 1935
8. Bradfield, Richard
"The value and limitations of calcium in soil
structure."
Am. Soil Survey Assn. Rpt. 17:31-2 1936
9. Buehrer, T. F.
"The movement of gases through the soil as a
criterion of soil structure."
Arizona Agri. Exp. Sta. Tech. Bull. No. 39 1932
10. Daniel, H. A.
"The physical changes in soils of the Southern
High Plains due to cropping and wind erosion and
the relation between the sand + silt ratios in
clay
these soils."
Journ. Am. Soc. Agron. 28:570-80 1936

11. Daniel, H. A., and Langham, W. H.
 "The effect of wind erosion and cultivation on the total nitrogen and organic matter content of soils in the Southern High Plains."
Jour. Am. Soc. Agron. 28:587-96 1936
12. Dean, H. L., and Walker, P. H.
 "A comparison of glass and quinhydrone electrodes for determining pH of some Iowa soils. II. The variability of results."
Jour. Am. Soc. Agron. 27:519-25 1935
13. DeTurk, E. E.
 "Soil conservation from the viewpoint of soil chemistry."
Jour. Am. Soc. Agron. 29:93-112 1937
14. DeTurk, E. E., Bauer, L. C., and Smith, L. H.
 "Lessons from the Morrow Plots."
Ill. Agric. Exp. Sta. Bull. 300 1927
15. Dorman, Clarence
 "A comparative study of cropped and virgin soils."
Soil Science 36:101-17 1933
16. Machlis, J. A., and Williams, B. H.
 "Soil Survey of Hyde County, South Dakota." 1925
17. Millar, C. E.
 "Studies on virgin and depleted soils."
Social Science 16:433-37 1923
18. Rhoades, H. F.
 "Aggregate analysis as an aid in soil structure studies."
Am. Soil Survey Assn. Rpt. 13:165-9 1932
19. Shedd, O. M.
 "A comparison of the calcium content of some virgin and cultivated soils of Kentucky by an improved method for estimation of the element."
Ky. Agri. Exp. Sta. Res. Bull. 236 1921
20. Swanson, C. O.
 "The loss of nitrogen and organic matter in cultivated Kansas soils and the effect of this loss on the crop producing power of the soil."
Jour. Ind. and Eng. Chem. 7:529-32 1915
21. Swanson, C. O., and Miller, R. W.
 "The sulfur content of some typical Kansas soils, and the loss of sulfur due to cultivation."
Social Science 3:139-48 1917

22. Welker, R. H., Brown, P. E., and Young, A. W.
"Some chemical and bacteriological effects of
various kinds and amounts of lime on certain
southern Iowa soils."
Is. Agrl. Exp. Sta. Bull. 148 1932
23. Watkins, W. I., Deardorff, C. E., Hutton, J. G., and
Machlis, J. A.
"Soil Survey of Beadle County, South Dakota."
1924
24. Watkins, W. I., and Larson, G. A.
"Soil Survey of Moody County, South Dakota." 1926
25. Wilson, B. D.
"Exchangeable calcium and potassium in soils
as affected by cropping and fertilization."
Soil Science 29:91-100 1930
26. Wright, C. H.
Soil Analysis
(Thos. Murby and Co.) 1934
27. Yoder, R. E.
"A direct method of aggregate analysis of soils
and a study of the physical nature of erosion
losses."
Am. Soil Survey Assn. Rpt. 17:165 1936